

A young woman with short, straight blonde hair and bangs is shown in profile, looking out a window. She is wearing a red top. The background is a blurred view of greenery outside the window. The overall mood is contemplative and serene.

ORYZON

Pioneering personalized
medicine with **epigenetics**

Corporate Presentation
1Q 2026
ORY:SM / ORY.MC

Legal Notice

DISCLAIMER This document has been prepared by Oryzon Genomics, S.A. exclusively for informational purposes and to be used during the presentation. As a consequence thereof, this document may not, without the express and prior written consent of Oryzon Genomics, S.A., be disclosed or published, nor used by any other person or entity, for any other reason. Neither Oryzon Genomics, S.A., nor any entity acting on its behalf, assumes liability for this document if it is used with a purpose other than the above. The information and any opinions or statements made in this document have not been verified by independent third parties; therefore, no express or implied warranty is made as to the impartiality, accuracy, completeness or correctness of the information or the opinions or statements expressed herein. Neither Oryzon Genomics, S.A., nor any entity acting on its behalf, assumes liability of any kind, whether for negligence or any other reason, for any damage or loss arising from any use of this document or its contents. Neither this document nor any part of it constitutes a contract, nor may it be used for incorporation into or construction of any contract or agreement. After careful review, Oryzon Genomics, S.A. considers that the information contained herein does not constitute “inside information” (as defined in Article 7 of Regulation (EU) No 596/2014 of the European Parliament and of the Council of 16 April 2014 on market abuse). In any case, this presentation should be read in conjunction with and taking into account all publicly available information of Oryzon Genomics, S.A., including the registration document approved and registered by the Comisión Nacional del Mercado de Valores (CNMV). The financial information contained herein is reflected in or has been extracted from Oryzon Genomics, S.A.’s public financial statements, and shall be qualified in its entirety by the information contained in such financial statements, copies of which are available on the websites of Oryzon Genomics, S.A. (www.oryzon.com) and the CNMV (www.cnmv.es). This document may contain summarized information or information that has not been audited, and certain financial and statistical information in this presentation has been subject to rounding adjustments and to currency conversion adjustments. Accordingly, the sum of certain data may not conform to the expressed total.

IMPORTANT INFORMATION This document does not constitute an offer or invitation to purchase or subscribe shares in accordance with the provisions of Regulation (EU) 2017/1129 of the European Parliament and of the Council of 14 June 2017, and/or the restated text of the Securities Market Law, approved by Law 6/2023 of 17 March, and its implementing regulations. Nothing in this document constitutes investment advice. In addition, this document does not constitute an offer of purchase, sale or exchange, nor a request for an offer of purchase, sale or exchange of securities, nor a request for any vote or approval in any jurisdiction. The shares of Oryzon Genomics, S.A. may not be offered or sold in the United States of America except pursuant to an effective registration statement under the Securities Act of 1933 or pursuant to a valid exemption from registration.

FORWARD-LOOKING STATEMENTS This communication may contain forward-looking information and statements about Oryzon Genomics, S.A., including financial projections and estimates and their underlying assumptions, statements regarding plans, objectives and expectations with respect to future operations, capital expenditures, synergies, products and services, and statements regarding future performance. Forward-looking statements are statements that are not historical facts and are generally identified by the words “expects,” “anticipates,” “believes,” “intends,” “estimates” and similar expressions. Although Oryzon Genomics, S.A. believes that the expectations reflected in such forward-looking statements are reasonable, investors and holders of Oryzon Genomics, S.A. shares are cautioned that forward-looking information and statements are subject to various risks and uncertainties, many of which are difficult to predict and generally beyond the control of Oryzon Genomics, S.A., that could cause actual results and developments to differ materially from those expressed in, or implied or projected by, the forward-looking information and statements. These risks and uncertainties include those discussed or identified in the documents sent by Oryzon Genomics, S.A. to the CNMV, which are accessible to the public. Forward-looking statements are not guarantees of future performance. They have not been reviewed by the auditors of Oryzon Genomics, S.A. You are cautioned not to place undue reliance on the forward-looking statements, which speak only as of the date they were made. All subsequent oral or written forward-looking statements attributable to Oryzon Genomics, S.A. or any of its members, directors, officers, employees or any persons acting on its behalf are expressly qualified in their entirety by the cautionary statement above. All forward-looking statements included herein are based on information available to Oryzon Genomics, S.A. on the date hereof. Except as required by applicable law, neither Oryzon Genomics, S.A., nor any entity acting on its behalf, undertakes any obligation to publicly update or revise any forward-looking statements, whether as a result of new information, future events or otherwise. Information in this document, if applicable, about the price at which securities issued by Oryzon Genomics, S.A. have been bought or sold in the past or about the yield on securities issued by Oryzon Genomics, S.A. cannot be relied upon as a guide to future performance.

Oryzon Develops Potent and Selective Epigenetic Drugs



Epigenetics experts specializing in LSD1 Biology



Iadademstat: in Phase II. Potentially best-in-class LSD1i developed in multiple hematology and oncology indications, with **well-defined registrational pathways for potential accelerated approval**



Vafidemstat: Phase III ready. LSD1i CNS asset geared toward ameliorating aggression and agitation in borderline personality disorder. **Phase II in schizophrenia**



Experienced management team, Board of Directors and world-renowned clinical regulatory and industry experts



Robust financial position with > \$60m raised in 2025; cash runway through 1H 2027

Oryzon: a Specialist in Epigenetics Developing New Targeted Therapies in Oncology/Hematology and CNS

Developing potent and selective epigenetic drugs with well-defined registrational pathways

Iadademstat

- Strong clinical data in unfit 1L AML (100% ORR, 90% strict CRs)
- Leverages CRADA-NCI agreement; 6 of 7 ongoing oncology trials sponsored by the NCI or top-tier U.S. institutions
- Oncology and Hematology: Multiple Phase Ib and II trials in different indications
- Unfit 1L AML: Final data expected at ASH-2026
- Phase Ib in Sickle Cell Disease. Clinical PoC I 4Q2026

Vafidemstat

- Phase III-ready asset in aggression in Borderline Personality Disorder (BPD)
- Phase II in Schizophrenia. Expected readout in 2H2027
- Phase II in aggression in ASD to start in 2026




IADADEMSTAT

A Phase II LSD1 inhibitor
for oncological and
hematological diseases


LSD1 inhibition MoA in AML and Other Malignancies is Well Characterized and Understood

Mechanism of Action

- LSD1 is required for leukemic stem cell survival and blocking leukemic cell differentiation
- **Iadademstat prevents leukemic stem cell survival and promotes rapid differentiation/death of leukemia cells**



Iadademstat: First and Potentially Best-in-Class LSD1 Inhibitor for Hematological Diseases



The Asset

- The most potent (nM) oral LSD1 inhibitor in clinical development
- Safety profile well-established: +225 pts treated with iadademstat in multiple Phase I/II trials
- Orphan Drug Designation for AML (US and EU)



ARTICLE · Volume 33, Issue 3, P495-511.E12, March 12, 2018 · [Open Archive](#)

[Download Full Issue](#)

ORY-1001, a Potent and Selective Covalent KDM1A Inhibitor, for the Treatment of Acute Leukemia

[Tamara Maes](#)^{1,6} [Cristina Mascaró](#)¹ · [Iñigo Tirapu](#)¹ · ... · [Matthew Fyfe](#)¹ · [Julio Cesar Castro-Palomino](#)¹ · [Carlos Buesa](#)¹ ... [Show more](#)



First-in-Human Phase I Study of Iadademstat (ORY-1001): A First-in-Class Lysine-Specific Histone Demethylase 1A Inhibitor, in Relapsed or Refractory Acute Myeloid Leukemia

Authors: [Olga Salamero, MD](#)¹ · [Pau Montesinos, MD](#)² · [Christophe Willekens, MD](#)³ · [José Antonio Pérez-Simón, MD, PhD](#)⁴ · [Arnaud Pigneux, MD, PhD](#)⁵ · [Christian Récher, MD, PhD](#)⁶ · [Rakesh Popat, MB, BS, PhD](#)⁷ · ... [SHOW ALL](#) ... and [Tim C. P. Somervaille, MBBS, PhD](#)⁸ | [AUTHORS INFO &](#)

ARTICLES · Volume 11, Issue 7, E487-E498, July 2024

[Download Full Issue](#)

Iadademstat in combination with azacitidine in patients with newly diagnosed acute myeloid leukaemia (ALICE): an open-label, phase 2a dose-finding study

[Olga Salamero, MD](#)^a · [Antonieta Molero, MD](#)^a · [José Antonio Pérez-Simón, MD](#)^b · [Montserrat Arnan, MD](#)^c · [Rosa Coll, MD](#)^d · [Sara García-Avila, MD](#)^e et al. [Show more](#)




Iadademstat in Oncology and Hematology: Multiple Opportunities Leveraging CRADA-NCI Agreement




Indication	Sponsor	Preclinical	Phase I	Phase II	Phase III	Status/Upcoming catalysts*
Acute Myeloid Leukemia (AML) 1L unfit patients: combination w/ azacitidine	Oryzon	ALICE				Completed. Published (Lancet Hematol)
1L AML unfit patients: combination w/ azacitidine + venetoclax	OHSU	IIS-ALICE-2				EHA-2026/ASH-2026
1L AML unfit patients: combination w/ azacitidine + venetoclax	NCI	CRADA-AML				
Refractory/Relapsed AML FLT3 mutation+ pts, combination w/ gilteritinib	Oryzon	FRIDA				EHA-2026/ASH-2026
Myelodysplastic Syndrome (MDS) combination w/ azacitidine	MCW	IIS-X005				ASH-2026
MPN: combination w/ ASTX727	NCI	CRADA-MPN				ASH-2026
Extensive-Disease Small Cell Lung Cancer (ED-SCLC) 1L patients: combination w/ ICI	NCI	CRADA-SCLC				ESMO 2026
Extensive-Disease Small Cell Lung Cancer (ED-SCLC) 1L/2L pts: combination w/ ICI + SBRT	Yale	IIS-TIARA				
Sickle Cell Disease (SCD)	Oryzon	RESTORE				ASH-2026
Essential Thrombocythemia (ET)	Oryzon	IDEAL				Approved by EMA. ASH-2026

Iadademstat, a pipeline in a single asset

Iadademstat Combinations in AML are Highly Encouraging. 100% ORR and 90% pure CR in 1L AML



Triple Combo Iada-Ven-Aza in First Line AML




ALICE-2

C. Lachowicz et al ASH 2025




ALICE-2



Adding Iadademstat to Standard of Care Provides the Most Comprehensive Leukemia Coverage

Addresses all three Key Drivers of Leukemia Versus Other Therapeutic Approaches

In intermediate- and high-risk patients, iadademstat provides a clear MoA advantage over the emerging gilteritinib triplet by targeting in addition the leukemic stem cell compartment



Iadademstat Approval Strategy in 1L AML by stratifying patients

- **ALICE-2:** Final data readout expected in **Q4 2026** (24 pts, for 21 evaluable pts).

Registrational ALICE-3: Seamless Phase II–III trial planned in adverse risk patients, with:

Accelerated Approval (AA) based on **CR-based endpoints**, and

Full approval based on **overall survival (OS)**.


Trial initiation expected in **2027**.

ALICE-3 accrual timeline:

Expected to require approximately **~24 months** for full enrollment

Accelerated approval strategy guided by previous combination trial designs approved by FDA, supporting the regulatory pathway and selected endpoints



LSD1 Inhibition and ladademstat Show Activity in the Whole Spectrum of Myeloproliferative Diseases



ladademstat's direct activity against leukemic stem cells, together with its differentiation-inducing and pro-apoptotic effects, **supports its potential role as a backbone therapy across multiple combination regimens throughout the disease spectrum**

Small Cell Lung Cancer (SCLC): ladademstat and anti-PD-L1 Combination Inhibits SCLC Progression in Preclinical Model

- ladademstat renders the SCLC cells visible to the immune system
- ladademstat synergizes with ICIs and boosts the host immune system by increasing T cell infiltration and preventing T-cell exhaustion



Analysis of epigenetic determinants of antigen presentation identified LSD1 gene expression as a correlate of worse survival outcomes for patients treated with either nivolumab or the combination of nivolumab and ipilimumab

Ongoing Phase I/II SCLC Trials in Combination with ICI

A Phase I Dose Finding and Phase II Randomized Trial of Iadademstat Combined With Immune Checkpoint Inhibition Maintenance After Initial Chemoimmunotherapy in Patients With Extensive-Stage Small Cell Lung Cancer

CRADA-SCLC (NCT06287775)

Sponsor: National Cancer Institute

PI: Dr. Charles Rudin (MSKCC)



Enrollment (Estimated)

45 pts

Primary Objective

To compare the progression-free survival (PFS) between the combination of iadademstat plus ICI versus ICI maintenance alone.

Secondary Objectives

- To compare objective response rate (ORR) and overall survival (OS) between treatment arms.
- To evaluate the safety of combination iadademstat plus ICI.

RECRUITING (More than 30 sites across the U.S.)

Iadademstat and Radiation Therapy With Atezolizumab in Extensive Stage Small-cell Lung Cancer (ES-SCLC) Patients With Persistent, Recurrent or Progressive Disease After First Line Systemic Therapy

IIS-TIARA (NCT07113691)

Sponsor: Yale University

PI: Dr. Anne Chiang (Yale University)

Enrollment (Estimated)

15 pts

Primary Objective

To evaluate the safety of iadademstat combined with atezolizumab and stereotactic body radiotherapy (SBRT)

Secondary Objectives

To evaluate the efficacy of iadademstat combined with atezolizumab and SBRT in terms of disease control rate (CR, PR, SD), ORR (CR or PR), duration of response, progression-free survival, OS, and local control of irradiated target lesion

RECRUITING



**Non oncological
hematological disorders:
SCD and ET**


Sickle Cell Disease

Caused by a mutation in B-chain of Hemoglobin (Hb)

Around 20-25 million people are living with SCD across the globe

Sickle Cell Disease: High Unmet Medical Need

- Severe, lifelong genetic disorder with significant morbidity and early mortality
- **HSCT is the only curative option**, accessible to few patients
- Urgent need for **targeted, disease-modifying therapies** to improve survival and quality of life
- Key strategies: **increase fetal hemoglobin (HbF)** and **reduce hemolysis**



Strong Activity and High Interest from Leading Pharma Companies

USA average annual direct healthcare costs per adult patient year is > \$100,000, Annual US healthcare costs are > \$2 B




- 2019: Oxbryta received accelerated FDA approval
- 2022: Oxbryta achieved \$328 million in U.S. sales
- Pfizer demonstrated in 2yr a significant market opportunity in sickle cell disease (SCD)



**Global Addressable Patient
Population In Developed Countries
~320,000**

Iadademstat MoA in Sickle Cell Disease: An FDA-approved MoA

LSD1 is a component of the protein complexes that repress *HBG1/2* transcription. Iadademstat may restore *HBG1/2* expression by inhibiting these repressive complexes



Strong Preclinical Data

- In rodents
- In Baboons in single dose
- In Baboons in long term dosing
- In ex-vivo human blood

Modified from:

- Suzuki et al.. *Fetal globin gene repressors as drug targets for molecular therapies to treat the β-globinopathies*. Molecular and Cellular Biology. 2014 Oct;34(19):3560-3569. DOI: 10.1128/mcb.00714-14.
- Paikari, A., Sheehan, V. *Fetal haemoglobin induction in sickle cell disease*. Br J Haematol 2018, 180(2): 189-200. DOI 10.1111/bjh.15021

An Open-label Phase Ib in SCD (RESTORE Trial) Approved by EMA

First two cohorts enrolled; PoC clinical activity expected in 2026

RESTORE

Sponsor: Oryzon

Size N~24-30:

- Escalation up to n=18
- Expansion n=12

Dose escalation, followed by dose expansion at RP2D

Open-label, up to 24 weeks treatment

Endpoints

Primary:

Safety and tolerability, RP2D

Secondary:

HbF induction, PK/PD, hemolysis markers

Exploratory:

Vaso-occlusive crisis frequency and duration, effect on transfusion, PROs, pharmacogenomics

Status

- ✓ Recruiting
- ✓ **By 1H 2026**, we expect to establish safety in this population and obtain initial Biomarker (HbF) data as PoC of MoA
- ✓ **By 2H 2026**, we expect to have a first assessment of therapeutic efficacy (HbF) in SCD

Iadademstat Approval Strategy in Hematology

- **RESTORE:** Final data readout expected in **Q2 2027 (24-30 patients)**.

- **Registrational RESTORE-2:** Seamless Phase II–III trial planned, with:

- **90+45 pts → AA** based on **HbF-based endpoint**, and
- **Full approval** based on **VOCs**.
- Trial initiation expected in **2-3Q 2027**.

- **RESTORE-2 accrual timeline:**

Expected to require approximately **18 months** for full enrollment.

Essential Thrombocythemia (ET): A Fast Follower Strategy

Essential Thrombocythemia (ET) Overview

- ET is the most common type of myeloproliferative neoplasm (MPN) and increases the risk of serious complications such as stroke, heart attack, and pulmonary embolism.
- The disease affects **~200,000 people in the U.S.**

Mechanism of Action


- LSD1 inhibition blocks the terminal differentiation of megakaryocytes into platelets, leading to a steady reduction in circulating platelet counts.

Competitive Landscape

- MSD published positive Phase II results with bomedemstat, another LSD1i
- MSD initiated a Phase III trial (Shorespan-007) in 2024, with results expected in 2027

Iadademstat Positioning

- Iadademstat is the only advanced LSD1 inhibitor in development across oncology and hematology (Phase Ib and II), with +225 patients treated.
- A Phase II clinical trial (IDEAL) has been approved by EMA and is planned to start in 1Q2026 to provide PoC.



**ORYZON developing
first-in-class epigenetic
drugs in CNS**

VAFIDEMSTAT

**A Phase III-ready LSD1 inhibitor for CNS
diseases**


LSD1, an epigenetic key enzyme in CNS

It plays a critical role in neuro-genesis and the regulation of cortical development

It localizes in-vivo to enhancers and promoters of confirmed CNS disease risk genes

It has been involved in neuro-developmental diseases

ORYZON is the only company to have developed an LSD1i for CNS: vafidemstat



Vafidemstat pharmacology supports use in different mental diseases

Vafidemstat (aka ORY-2001) and other LSD1i induce expression of genes **involved in neuronal plasticity**, restoring neuronal morphology, branching and axonal navigation

Vafidemstat **restores the response to stress** by regulating genes involved in control of stress cues in the PFC-amygdala axis, as IEG, SRF, and others

LSD1i is able to **rescue glutamatergic NMDA-R hypofunction** in prefrontal cortex in different ASD and SCZ models

Vafidemstat improves sociability

Vafidemstat reduces aggression

Vafidemstat improves memory

Borderline Personality Disorder, Schizophrenia, Autism, ADHD, others

Vafidemstat is Safe and Well Tolerated

A very robust safety package. +425 treated subjects



Oral & Brain Penetrant

Oral, once daily
1.2 mg /day (RP2D)

An optimal CSF: plasma
ratio of 0.9



Safe, No DDIs

Comparable SARs between
placebo and vafidemstat arms
in 6 Phase II trials



No side effects

No weight gain
No sedation / somnolence
No sexual dysfunction
No extrapyramidal signs


Aggression, a Huge Medical Need in CNS Diseases



Agitation and aggression are common in most psychiatric and neurodegenerative diseases



Clinical data with vafidemstat



Vafidemstat Current Clinical Development

- Exploring large multifactorial indications (Borderline Personality Disorder, Schizophrenia and Autism)
- Exploring also feasibility in some rare genetically-driven neurodevelopmental disorders (Phelan McDermid, Fragile X, Kabuki, etc)

Indication	Sponsor	Preclinical	Phase I	Phase II	Phase III	Status/upcoming catalysts
Borderline Personality Disorder (BPD) Agitation/Aggression	Oryzon	PORTICO-2			Submitted	Phase III in preparation
Schizophrenia Negative Symptoms / Positive Symptoms / CIAS	Oryzon	EVOLUTION				EU expansion in 2026; readout in 2H2027
Autism Spectrum Disorder (ASD) Aggression / Repetitive Behavior	Oryzon	HOPE-2				PhII in preparation; to initiate in 1H2026

Vafidemstat Demonstrated a Relevant Clinical Benefit in Reducing Agitation /Aggression across ASD, ADHD and BPD Patients in PoC Phase IIa Study

PCN Psychiatry and Clinical Neurosciences



Regular Article | [Open Access](#) |


REIMAGINE: A central nervous system basket trial showing safety and efficacy of vafidemstat on aggression in different psychiatric disorders

Marc Ferrer MD, PhD, Vanesa Richarte MD, PhD, Laura Gisbert MD, PhD, Jordi Xaus PhD, Sonia Gutierrez BSc, MSc, Maria Isabel Arevalo PhD, Michael Ropacki MA, PhD, Roger Bullock MD, Carlos Buesa PhD , Josep Antoni Ramos-Quiroga MD, PhD


First published: 12 February 2025 | <https://doi.org/10.1111/pcn.13800>

Clinical Trial Registration: REIMAGINE EudraCT#: 2018-002140-88.

Eight-week vafidemstat treatment led to a statistically significant reduction in agitation/aggression compared with baseline across all assessments (all participants, $p < 0.0001$)



Borderline Personality Disorder: an Unmet Medical Need & Vast Commercial Opportunity (+\$3Bn)



Prevalent & impairing disease

9 million in US & EU

Two main types of symptoms

Psychiatric symptoms
+
Agitation/Aggression
(including self-aggression)

No approved drugs yet

Patients on off-label anti-psychotics

Vafi improves these symptoms in:

- BPD patients
- PC models

Oryzon is leading the BPD field ahead of the competition

PORTICO: a Global Phase IIb Randomized, Placebo-Controlled, Double Blinded Trial in BPD to Inform Subsequent Development

Key inclusion criteria

Men and women 18-65 years of age

DSM-5 BPD diagnostic criteria, at least 3 months before the Screening visit.

Agitation-Aggression Psychiatric Inventory-Clinician Report (AAPI-CR) Agitation & Aggression (A/A) subscale score of ≥ 16 (severity x frequency) summed across the 4-items comprising the A/A subscale, and the sum of the A/A subscale severity scores ≥ 6

Stable regimen of background pharmacotherapy at Screening, Baseline and throughout the trial

Maintenance of pre-screening psychotherapy schedule throughout the trial

Willing and able to adhere to the protocol prohibitions, restrictions and requirements

N=211
Randomized
1:1

Vafidemstat, 1.2mg
Once daily (5 ON, 2 PBO), N=106

Placebo
Once daily, N=105

14-week trial

Endpoints

Primary:

Agitation/Aggression (CGI-S A/A) from baseline to weeks 8-12

Improvement in Borderline Personality Disorder Checklist (BPDCL) from baseline to weeks 8-12

Secondary (efficacy):

To evaluate the change over time on the CGI-S A/A

To evaluate the change over time on the BPDCL


To evaluate the difference on the following measures, from baseline to weeks 8-12, as well as change over time, between the active treatment arm and the placebo arm:

- ❖ Borderline Evaluation of Severity over Time (BEST)
- ❖ State-Trait Anger Expression Inventory 2 (STAXI-2)
- ❖ State-Trait Anxiety Inventory (STAI)
- ❖ Beck Depression Inventory – II (BDI-II)


PORTICO final results presented at **ECNP-2024**

PORTICO: Vafidemstat Improves Aggression Over Placebo (Secondary Endpoint)


“STAXI allows us to effectively and efficiently know if someone is getting better over the course of the trial and in ways that are highly relevant to the clinical work with BPD” - Sarah Fineberg, M.D., Ph.D.



Primary endpoint CGI-S A/A (Clinician rated)



Secondary endpoint STAXI-2 Trait Anger (Patient rated)



PORTICO: Vafidemstat was Safe and Well Tolerated

Vafidemstat-treated patients showed a reduced inclination towards self-harm

Treatment-Emergent Adverse Events by Preferred Term Occurring in > 5% of Subjects	Placebo	Vafidemstat
	(N=104)	(N=106)
	N (%), e	N (%), e
TEAEs by Preferred Term	68 (65.4%), 214	61 (57.5%), 192
Headache	17 (16.3%), 18	13 (12.3%), 16
Nasopharyngitis	18 (17.3%), 22	9 (8.5%), 11
Tension Headache	6 (5.8%), 17	5 (4.7%), 11
Platelet Count Decreased	1 (1.0%), 1	8 (7.5%), 8*
Nausea	2 (1.9%), 2	6 (5.7%), 6
Intentional Self-Injury	6 (5.8%), 10	1 (0.9%), 2

Vafidemstat: Summary and Next Steps

End-of-Phase II Meeting with FDA resulted in positive feedback:

- Agitation-Aggression in BPD acknowledged as a possible therapeutic indication
- FDA feedback supports initiation of a Phase III trial (PORTICO-2) using STAXI-2 Trait Anger as a primary efficacy endpoint measure, but additional information was requested regarding the face validity of the proposed endpoints.
- Secondary endpoints will include patient-rated and clinician-rated scales to assess agitation/aggression and overall BPD improvement

Phase III Protocol submitted after further constructive interactions with the agency:

- In response, the company convened a panel of renowned U.S. experts (including Dr Alan F. Schatzberg, Dr. Eric Hollander, Dr Emil F. Coccaro and Dr Sarah Finneberg) to contribute to the design of the Phase III protocol.
- As per FDA suggestions, the initial design was modified to incorporate a key secondary endpoint, the Overt Aggression Scale-Modified, OAS-M, a well-validated ClinRO shown to reliably measure reductions in aggression in patients receiving pharmacological treatment. Importantly, this scale has a strong psychometric correlation with the proposed primary endpoint, the STAXI-2 Trait Anger scale.
- Phase III protocol also included Qualitative research and Psychometric analyses following FDA recommendations
- **FDA's feedback received on Oct 16th**, including guidance on study endpoints and certain non-clinical considerations

Next Step:

Addressing FDA's comments; protocol resubmission targeted before year-end

New and Prestigious US-centric Clinical Advisory Board for CNS



Alan F. Schatzberg



Alan F. Schatzberg renowned American psychiatrist. Since 1991, he has been the Kenneth T. Norris Jr. Professor of Psychiatry and Behavioral Sciences at Stanford University School of Medicine. He was chair of the department Psychiatry and Behavioral Sciences at Stanford from 1991 to 2010. He is also the co-editor-in-chief of the Journal of Psychiatric Research. Alan Schatzberg, was the principal investigator for mifepristone for use as an antidepressant developed by Corcept Therapeutics, a company Schatzberg had founded.



Dr. Emil F. Coccaro



Dr. Emil F. Coccaro is a psychiatrist in Columbus, Ohio and is affiliated with Ohio State University Wexner Medical Center. He received his medical degree from New York University Grossman School of Medicine and has been in practice for more than 20 years. He is an expert in Aggression and has contributed to The Overt Aggression Scale Modified (OAS-M) for clinical trials targeting impulsive aggression and intermittent explosive disorder.




Eric Hollander, M.D.



Eric Hollander, M.D. is Professor, Department of Psychiatry and Behavioral Sciences at Albert Einstein College of Medicine in NYC. Director, of the Autism and Obsessive Compulsive Spectrum Program, Department of Psychiatry and Behavioral Sciences.




Dr. Sarah Fineberg



Dr. Sarah Fineberg is Assistant Professor of Psychiatry at Yale University investigating the neurobiological mechanisms behind borderline personality disorder (BPD) and related mental health conditions. She has participated in several BPD clinical trials.

Vafidemstat: PORTICO-2 Phase III Study Proposed Design

Aggression: Primary endpoint (STAXI-2 Trait Anger) + Key Secondary endpoint (OAS-M)



Endpoints

Primary:

Efficacy in Agitation-Agression by STAXI-2 Trait Anger

Key Secondary:

Efficacy in Agitation-Agression by OAS-M (ClinRO FDA accepted scale)

Secondary:

Efficacy improvements in:

- Overall improvement by BEST
- Overall improvement by CGI-S
- Depression by BDI-II

Safety

Exploratory:

- PK
- Target Engagement
- Exploratory biomarkers
- Genetic Polymorphisms

Expanding Aggression Program: New Trials Under Preparation in ASD

To be conducted initially in Spain financed through the EU-IPCEI grant (13.1 Million Euros)



Phelan-McDermid Syndrome (PMS)

A rare genetic condition (prevalence estimate 1/30,000 births) primarily caused by a terminal deletion on chromosome 22q13.

Aggressive behavior is seen in approximately 25% of affected individuals.

Idiopathic ASD

Up to 68% of ASD patients may show agitation & aggression

Aggression is one of the reasons why young patients with ASD are institutionalized as adults

Exploring strategic collaborations with biotech companies to leverage AI platforms to select patients most likely to benefit from vafidemstat

Schizophrenia Still an Enormous Unmet Medical Need

Despite the approval of Cobenfy (BMS) for positive symptoms, addressing negative or cognitive symptoms, as well as treatment-resistant schizophrenia, remains a significant challenge in managing this disease

A prevalent & impairing disease
20 million ww.



~5 million in
US & EU

Three main types of symptoms



Positive or Negative +
Cognitive Impairment

No approved drugs yet for Negative
Symptoms or Cognitive Impairment



Vafi improves these
symptoms in PC models

Total Addressable Market
in 2024



US\$ +10
billion

Highest Revenue Drug Category
long-acting injectable (LAI)
antipsychotics



Single Best seller: + \$4.1 Billion
Cobenfy expected peak sales + 6 Bn

Moderate
competition



Vafidemstat in Schizophrenia (SCZ)



Genetic link
between LSD1 and
SCZ



Preclinical in-vitro
and animal models
data supporting
LSD1 inhibition as a
new MoA in SCZ



No approved drugs
yet in negative
symptoms or
Cognitive
Impairment
symptoms



Strong market
interest in
Schizophrenia

EVOLUTION Phase IIb – a Real-World Trial

- **EVOLUTION** is designed as a **real-world trial**, with inclusion and exclusion criteria crafted to permit comorbidities, provided they are stable and/or under treatment.
 - Participants must present with predominant negative symptoms (PNS) of schizophrenia.
 - All enrolled subjects must be on stable treatment for schizophrenia, receiving no more than one atypical antipsychotic to manage psychotic symptoms. Long-acting injectable (LAI) formulations are permitted. The concomitant use of a second antipsychotic is not allowed.
 - 24 weeks of treatment to assess possible variation on CIAS
- A high-quality trial designed to meet the standards of registrational studies in the indication: multicenter, double-blind against a control arm, with regulatory-accepted endpoints

EVOLUTION: A Phase II Study to Measure Negative Symptoms in SCZ

Recruitment ongoing in Spain. Trial expansion unfolding in Eastern Europe countries

Endpoints

Primary:

Improvement in Negative Symptoms


Secondary - Efficacy

Improvements in:

- Cognitive impairment associated with SCZ (CIAS)
- Positive symptoms for SCZ
- Positive and Negative Syndrome Scale (PANSS) Total Score
- Functional impairment in adult SCZ patients

Secondary - Safety:

To evaluate vafidemstat safety in adult SCZ patients



Readout: expected in 2027




ORY-4001

A selective HDAC6 inhibitor
for ALS, CMT and other
CNS diseases

ORY-4001 is a highly potent and selective HDAC6 inhibitor

HDAC6 has been suggested as a therapeutic target for Amyotrophic Lateral Sclerosis (ALS), Charcot-Marie-Tooth (CMT), and other CNS diseases

- **Highly potent and selective** HDAC6 inhibitor with good pharmacology
- **Efficacy in a CMT1A model:**
 - increases nerve conduction velocity and CMAP
 - increases axonal number and myelination
- **Efficacy in ALS models** in mice, zebrafish, nematodes, and ALS patient-derived lymphoblasts
- 0.5M USD grant received from the U.S. ALS Association to support regulatory preclinical development in ALS
- First-in-Man readiness expected in 2H2026
- Available for partnering



Sacilotto N et al. ORY-4001, a novel potent and selective oxadiazole-based HDAC6 inhibitor shows pre-clinical therapeutic efficacy in CMT1A. PNS 2023 annual meeting

Oryzon Uniquely Positioned to Pioneer Epigenetic Drugs



Epigenetics experts specializing in LSD1 Biology



Iadademstat: LSD1i asset developed in multiple hematology and oncology indications, with potential accelerated approval pathways in 1L AML and SCD



Vafidemstat: Phase III ready LSD1 CNS asset geared toward ameliorating aggression and agitation



Experienced management team, Board of Directors and world-renowned clinical experts



Robust financial position; cash runway through 1H2027

Upcoming clinical development milestones

- **Clinical data updates in 1L AML** in combo w/ venetoclax/azacitidine:
 - Interim update at EHA-2026 (~75% of the planned recruitment)
 - **Final data expected at ASH-2026**
- **Additional clinical data updates in several other trials**
- **Sickle Cell Disease: safety and PoC clinical activity in 2026**
- **ET PhII** (EMA approved) to start in 1Q2026
- PhIII **Borderline Personality Disorder** protocol resubmission to FDA by year-end
- **Schizophrenia** EU expansion in 2026; readout expected in 2H2027
- **Autism Spectrum Disorder PhII** to initiate in 1H2026

A photograph of a modern glass skyscraper with the Oryzon logo on top. The logo consists of the word "ORYZON" in white capital letters on a dark rectangular background, followed by a square icon containing a stylized globe. The building's glass facade reflects the sky and clouds.

ORYZON

Pioneering personalized medicine in **epigenetics**